



Climate Change Vulnerability of Terrestrial Areas in Chesapeake & Ohio Canal National Historical Park - East

PLANNING FOR CHANGE

Located in a zone of ecological transition, spanning the Northern Piedmont from the Blue Ridge to the Coastal Plain, **Chesapeake & Ohio Canal National Historical Park - East (CHOH-E, C&O Canal - East)** supports a diversity of natural communities. As a 19,500-acre protected area within a landscape heavily pressured by development, the park provides opportunities for ecological adaptation in the face of environmental change. However, as global temperatures rise and precipitation patterns become more extreme (Intergovernmental Panel for Climate Change (IPCC) 2013), the *natural habitats* (see glossary, page 9, for definition of italicized words) of the park are increasingly vulnerable to climate-driven stresses. To begin to understand how climate change will impact terrestrial habitats in the National Capital Region (NCR), the National Park Service partnered with NatureServe to conduct **climate change vulnerability assessments**. Because the park is 184 miles long, we split it into two assessments: East and West. See the brief for C&O Canal - West to see results for the western portion of the park. This initial assessment for C&O Canal - East examines how climate and landscape influence vulnerability and identifies areas most and least vulnerable today and in the near future. Our goal is to help park managers and visitors understand climate change vulnerability and support the development of informed management strategies.



| IMPORTANT TERMS |
|--|
| VULNERABILITY Describes whether the ecological condition of a region or habitat is at risk from climate change impacts. |
| EXPOSURE Describes the nature and magnitude of changes in temperature and precipitation by comparing a given time period against a historical baseline. |
| ADAPTIVE CAPACITY Describes the ability of a region or habitat to maintain species and ecological processes as climate changes. Adaptive capacity at a specific location considers the area's unique physical features and its connectivity to other natural areas. |

C&O CANAL - EAST is located in the northeastern portion of the NCR, following the Potomac River from east of Harpers Ferry to Washington D. C. We consider the implications of climate change for the park within the context of the NCR surrounding area and the Mid-Atlantic region.


WHAT IS CLIMATE CHANGE VULNERABILITY?

We define **VULNERABILITY** as the risk of losing species and ecosystem processes due to rapid environmental change. We integrate two components of vulnerability, **EXPOSURE** and **ADAPTIVE CAPACITY**, to arrive at a single measure. Areas most at risk are those likely to experience big changes in temperature and precipitation (i.e., high exposure) but have little capacity to adapt (i.e., low adaptive capacity). To understand how the climate of C&O Canal - East is changing over time, we analyzed temperature and precipitation over two time periods: **observed** (1981 – 2014) and **near future** (through 2040), compared to a mid-century **baseline** (1948 – 1980). The results presented here are part of a broader regional assessment of vulnerability for 11 NCR parks and the surrounding National Capital Region, interpreted in the context of regional environmental change in the Mid-Atlantic. Our approach describes drivers of *climate change* vulnerability, how vulnerability is spatially distributed, and a framework for park managers to anticipate and address changes in the coming decades.



HOW VULNERABLE IS C&O CANAL - EAST?


C&O Canal - East stretches from D.C. across southern Maryland, within a highly urbanized and rapidly growing area of the Mid-Atlantic, but an area that also supports important biodiversity. The eastern portion of the park spans from the Blue Ridge, through the Northern Piedmont, all the way to the Southern Plains (i.e., Coastal Plain). These *ecoregions*, which represent provinces of unique physiography, soils, climate, and vegetation, together contain a diversity of natural communities. As the climate changes, **the park can play an important role as part of a network of refugia and stepping stones**, connecting natural communities and providing opportunities for adaptation. To better understand how management actions can protect park resources and enhance the ecological role of the park in a changing world, we first assess the overall **VULNERABILITY** of terrestrial habitats at C&O Canal - East by combining measurements of **EXPOSURE** and **ADAPTIVE CAPACITY**. Next, we characterize the drivers of **EXPOSURE** (changes in temperature and precipitation) and **ADAPTIVE CAPACITY** (landscape characteristics).



C&O CANAL - EAST IN CONTEXT. CHO-E passes primarily through the Northern Piedmont (beige), but also touches the Blue Ridge (blue) and Southeastern Plains (pink). Areas of natural vegetation cover are represented by darker shading on the map.


Research Highlights

VULNERABILITY SCORE




During the **observed** time period (1981 – 2014), we find vulnerability is low for most of C&O Canal - East, reflecting low to moderate climate change exposure for most of the area. Differences in vulnerability are primarily driven by differences in adaptive capacity across the landscape. Because of the park’s enduring features, and connectivity to other natural areas, adaptive capacity is average or above for much of the park, particularly when compared to the region as a whole. In the **near future** (through 2040), vulnerability is moderate to high for the park due to a sharp increase in exposure, though not as high as in portions of the surrounding area. Increased exposure is expected to lead to species turnover, but in areas with high adaptive capacity, the park is expected to continue to support healthy communities of diverse plants and animals into the future especially if connectivity to other natural lands is maintained.

Observed Climate Change Vulnerability (1981 – 2014)



OBSERVED VULNERABILITY. Vulnerability is low (blue) for 86% of CHO-E. 14% of the park has moderate vulnerability, and there are no areas of high or very high vulnerability (red). Higher vulnerability occurs where adaptive capacity, especially connectivity, is low, such as in areas along roads or near adjacent urban expansion.

Near Future Climate Change Vulnerability (through 2040)




NEAR FUTURE VULNERABILITY. In the future, vulnerability increases dramatically due to increased exposure with moderate to high values (yellow and orange) recorded for CHO-E. However, high adaptive capacity means CHO-E is less vulnerable than the NCR surrounding area. No areas (0%) of the park have very high vulnerability scores (red).

COMPONENTS OF VULNERABILITY

Exposure

To understand how global climate change impacts C&O Canal - East, we analyze **EXPOSURE**, defined as the **nature and magnitude of changes in patterns of temperature and precipitation**. Analyses of annual and seasonal climate data reveal that the park is already experiencing climate change and indicate that park managers can anticipate significantly more change in the near future.

EXPOSURE is measured by analyzing annual and seasonal changes in temperature and precipitation over three time periods. We first characterize a mid-20th century **baseline** (1948 – 1980) and quantify its natural variability. We then compare **observed** climate (1981 – 2014), and **near future** climate (through 2040) to baseline conditions and measure the changes. Baseline and observed climate data is derived from weather station measurements, which reduces uncertainty. Near future projections are derived from an ensemble of 15 global climate models from the IPCC 5th Assessment Report (Taylor et al. 2012, IPCC 2013). Full methods will be documented in a technical report available on the NPS IRMA portal after project completion.




CHANGES IN TEMPERATURE. Between the baseline and observed periods, an annual temperature increase of 0.7°C (1.3°F) has already occurred at CHOH-E, pushing the mean observed temperature up to 13.3°C (56°F). Continued increases are expected in the near future, especially in summer where the average temperature is 2°C (3.6°F) higher than that of the baseline. These increases in near future temperature are a statistically significant departure from baseline conditions, as indicated by greater saturation in color (dark red).

CHANGES IN PRECIPITATION. In the observed period, slight increases in precipitation were recorded across both seasons. In the near future, that pattern is projected to change with decreases in precipitation for the wettest quarter (summer) and increases for the driest (winter). Greater saturation in color represents values that are more statistically significant. The most notable change is the near future increase in winter precipitation which represents a statistically significant departure from baseline conditions (dark green).

Research Highlights

At C&O Canal - East, temperatures in the **observed** period (green) already show evidence of warming when compared to the baseline. Modest increases in both summer and winter precipitation are detected.

In the **near future** (purple), we anticipate shifts to warmer and drier summers and warmer and wetter winters. Models indicate that by 2040, precipitation may decrease by close to 20% in the summer as temperatures rise 2°C (3.6°F). In the winter, precipitation is forecast to increase by nearly 50% while temperatures rise 1.5°C (2.7°F).




SEASONAL TEMPERATURE AND PRECIPITATION PATTERNS. Each dot represents the 30-year average for a sample point falling within in CHOH-E and the immediate surroundings.

COMPONENTS OF VULNERABILITY

Exposure

The Magnitude of Change




The magnitude of change refers to **how much EXPOSURE** C&O Canal - East has experienced in recent decades and how much is projected for the near future. By comparing the climate in the **observed** (1981 – 2014) time period to the historical variability in the **baseline** (1948 – 1980), we can understand whether temperature and precipitation patterns today are *typical* of the range of conditions experienced in the past. Temperature and precipitation values falling completely outside their historical range would indicate *novel* climatic conditions, to which ecosystems may not be adapted. The projected climate of the **near future** is assessed against the baseline in the same way. Analyzing and mapping EXPOSURE gives managers an understanding of the nature and magnitude of climate change, how it may vary across the landscape, and how exposure contributes to overall vulnerability.

MEASURING THE MAGNITUDE OF CHANGE. The summer temperature for CHOH-E has increased since the mid-century baseline. The average temperature for the observed period (solid red line) is higher than recorded average annual temperatures (dots) for almost all years in the baseline, falling nearly outside the range of historical variability (shaded gray bar). The average summer temperature for the near future (not shown) is projected to be over 25.2°C (78°F), and falls completely outside the range of historical variability.

Research Highlights


There is already evidence of changing climate in parts of the Mid-Atlantic region (pink), though observed climate averages do not fall completely outside the range of variation experienced in the past. Temperatures in the **observed** period have risen, but precipitation patterns are close to historical norms, and overall exposure for C&O Canal - East is relatively low (purple). In the **near future**, the Mid-Atlantic region and C&O Canal - East rapidly move towards high levels of exposure and novel climate conditions (yellow). The overall exposure score of 0.20 for the near future indicates that by 2040, conditions at the park are likely to be more extreme than in 80% of the baseline years. Extreme temperatures are the primary driver of the exposure score, though shifts in precipitation towards drier summers and wetter winters also contribute.



COMPONENTS OF VULNERABILITY

Adaptive Capacity


Adaptive Capacity is the ability of a region or habitat to maintain species and ecological processes as the climate changes. In this study, we focus on two key characteristics of adaptive capacity, **LANDSCAPE DIVERSITY** and **LOCAL CONNECTEDNESS** (Anderson et al. 2016), to assess C&O Canal - East and the terrestrial habitats within it. We identify places that are more or less resilient to the disturbances associated with climate change to help park managers develop informed adaptation strategies.



ADAPTIVE CAPACITY AT CHOH-E. On balance, adaptive capacity at CHOH-E is average to far above average when compared to similar settings in the region. Areas of higher adaptive capacity are found where a diversity of physical conditions is present, including Lock 28, Marble Quarry, Blockhouse Point, and Great Falls. Lower adaptive capacity is found in areas with far below average connectivity.

Landscape Diversity

Areas with high **LANDSCAPE DIVERSITY** contain a range of physical conditions and habitats that can support species as they adapt to a changing climate. Within a specific geophysical setting, landscape diversity is scored highest if there is a diversity of landforms (e.g., high ridges, steep slopes, coves, and floodplains), wide ranges in elevation, and a range of soil types. These *enduring features* are the foundation upon which ecological systems are built.



LANDSCAPE DIVERSITY BY ECOLOGICAL SYSTEM. For the ecological systems occurring at CHOH-E, the graph displays the proportion of area in each of 8 adaptive capacity categories based on landscape diversity. Dark colors indicate higher landscape diversity and thus higher adaptive capacity. Taller bars designate ecological systems that occupy a greater percentage of the park. Some of the more uncommon ecological systems (shortest bars) have the highest landscape diversity. Cove Forest and Mafic Glade and Barrens each occupy less than 1% of the park but have landscape diversity that is far above average (dark purple).

Research Highlights


Landscape diversity is average to far above average for most of C&O Canal - East. This is due to the high variety of topographic settings (e.g., ridgetops, steep slopes, coves, lower slopes, and floodplains) and relatively wide ranges in elevation. These areas within the park provide more microclimates when compared to similar settings within the region.

COMPONENTS OF VULNERABILITY

Adaptive Capacity

Local Connectedness

LOCAL CONNECTEDNESS measures the degree to which current land cover patterns (e.g., agriculture, forest, wetlands) are likely to support important ecological processes and the movement and dispersal of species. When natural areas are connected, opportunities are present for climate change adaptation, and vulnerability is reduced.



LOCAL CONNECTEDNESS BY ECOLOGICAL SYSTEM. For the ecological systems occurring at CHOH-E, we show the proportion of area in each category of adaptive capacity based on local connectedness. Dark colors indicate higher connectivity and thus higher adaptive capacity. Taller bars designate ecological systems that occupy a greater percentage of the park. Estimated connectedness of ecological systems at the C&O Canal – East is generally average to above (blue to purple). Some of the more uncommon ecosystems in the park have the highest local connectedness, while the most common ecosystem, the River Floodplain, scores the lowest compared to the rest of the park.

Research Highlights

Local connectedness is relatively high for most of C&O Canal - East. Most of the Potomac corridor from Turkey Run upstream is predicted to provide at least average connectivity. Some of the highest local connectedness scores in the eastern portion of the park are found in the vicinity of Blockhouse Point.




A pond near lock 15 along the C&O Canal in Chesapeake & Ohio Canal National Historical Park.

PATTERNS OF VULNERABILITY

Local Trends

Understanding how climate change (EXPOSURE) and local landscape characteristics (ADAPTIVE CAPACITY) influence vulnerability is a central goal of this assessment. We quantified and mapped these relationships for C&O Canal - East and the NCR surrounding area to identify what areas are most or least vulnerable today, and what we can expect in the near future. ADAPTIVE CAPACITY and EXPOSURE scores were divided into low, moderate, high, and very high categories and combined into the overall VULNERABILITY scores as presented on page 2 of this document and as illustrated below.



VULNERABILITY SCORE

LOW

- LOW EXPOSURE / HIGH ADAPTIVE CAPACITY
- LOW EXPOSURE / MODERATE ADAPTIVE CAPACITY
- MODERATE EXPOSURE / HIGH ADAPTIVE CAPACITY

MODERATE


- LOW EXPOSURE / LOW ADAPTIVE CAPACITY
- MODERATE EXPOSURE / MODERATE ADAPTIVE CAPACITY
- HIGH EXPOSURE / HIGH ADAPTIVE CAPACITY

HIGH


- MODERATE EXPOSURE / LOW ADAPTIVE CAPACITY
- HIGH EXPOSURE / MODERATE ADAPTIVE CAPACITY

VERY HIGH

- HIGH EXPOSURE / LOW ADAPTIVE CAPACITY



Observed Climate Change Vulnerability
CHOH-E
(1981 – 2014)



Near Future Climate Change Vulnerability
CHOH-E
(through 2040)


Research Highlights

In the **observed** period, vulnerability of C&O Canal - East is generally low, owing to generally moderate adaptive capacity and relatively low exposure. Some areas of the park are already classified as having moderate exposure, but these areas coincide with zones of moderate to high adaptive capacity. There, high landscape diversity and local connectedness along the Potomac River, provides some protective benefits in the form of habitat and stepping-stones for species movement and adaptation.

Exposure increases to high levels across the region in the **near future**, resulting in scores of moderate to high for C&O Canal - East. Differences in vulnerability are driven by differences in adaptive capacity. Vulnerability is lowest (moderate) in areas where above average landscape diversity coincides with large bands of connected forest along the Potomac River. Developing forward-looking adaptation strategies aimed at maintaining or increasing adaptive capacity will help ensure that the natural resources of C&O Canal - East can endure the stresses of rapidly changing conditions.

WHAT WE CAN DO

Recommendations & Management Strategies



NEAR FUTURE VULNERABILITY AT C&O CANAL - EAST

Mapping the spatial pattern of vulnerability can inform site-specific adaptation strategies. Developed areas in the regional map are dark gray. Vulnerability within park boundaries is seen in the inset.

This **climate change vulnerability assessment** analyzed several components of VULNERABILITY for C&O Canal - East and identified the areas that are most and least vulnerable and why. This work represents an important first step in the application of a climate-smart adaptation framework (Stein et al. 2014) through which park managers can identify, evaluate, and implement adaptation strategies suited to observed conditions and adaptable to conditions in the near future. Additional work is needed to refine vulnerability and adaptive capacity to include other direct and indirect impacts of a changing climate.

Broad adaptation strategies relevant to C&O Canal - East are provided in the Climate Change Vulnerability Impacts and Strategies table. These strategies can support managers as they begin planning for change. The next phases of the project will provide additional information to guide site-specific adaptation actions aligned with these strategies. This work includes assessment of ecosystem-specific drivers of vulnerability, identification of regional priorities for maintaining connectivity, and evaluation of field data to characterize the local impact of non-climate stressors on ADAPTIVE CAPACITY. Through these actions, we can safeguard the natural treasures of C&O Canal - East and ensure that they are resilient for future generations of park visitors.

CLIMATE CHANGE VULNERABILITY IMPACTS AND STRATEGIES TABLE

| VULNERABILITY SCORE | CLIMATE IMPACTS | STRATEGY |
|---------------------|--|---|
| Low | With high adaptive capacity and relatively low climate change exposure, these sites are least at risk. In the near future, there are no areas of low vulnerability at CHOH-E. | Manage for persistence , focus actions on preventing impacts by <i>non-climate stressors</i> (e.g., habitat loss, fragmentation, and invasive species). |
| Moderate | With moderate to high adaptive capacity and less exposure, these areas can continue to support diverse natural communities at CHOH-E. Moderate vulnerability areas, like Marble Quarry, Blockhouse Point, and Great Falls, provide opportunities for species to move and communities to adapt. | Encourage persistence but accommodate change . Actions should focus on (1) decreasing <i>non-climate stressors</i> to restore or enhance ecological integrity and (2) maintaining landscape connectivity to facilitate transitions. |
| High | Areas with high vulnerability in the near future have high exposure but moderate adaptive capacity. This includes much of the park. <i>Species turnover</i> and restructuring of communities at CHOH-E is likely. | Accommodate change and novel communities . Maintaining connected landscapes will support the persistence of diverse ecosystems, but actions should accommodate turnover of native species. Actions to maintain ecosystem functions and processes and limit biodiversity loss are favored (e.g., aggressive management of invasives, species translocations). |
| Very High | With high exposure and low adaptive capacity, areas with very high vulnerability may experience transformational changes likely to negatively impact overall biodiversity. Less than 1% of CHOH-E falls in this category in the near future. | Accommodate significant change and reevaluate management goals . Actions can be targeted for maintaining ecosystem functions and limiting biodiversity loss, but efforts aimed at maintaining existing ecological communities may not achieve desired outcomes. |

TO LEARN MORE

The information in this report is based on climate change scenarios from the Intergovernmental Panel on Climate Change (IPCC 2013), published climate datasets TopoWX (Oyler et al. 2014) and Prism (Daly et al. 2008), an analysis of terrestrial resilience for the eastern United States (Anderson et al. 2016), and analyses by NatureServe. To explore the data and learn more about our methods and the science behind climate change, please visit the links below.

1. Please visit the NCR Enduring Features Data Basin Gallery to further explore the information about Chesapeake & Ohio Canal National Historical Park presented in this brief. http://bit.ly/databasin_NCR
2. More information about the data and methods used to characterize exposure will be made available in the final project report (Smyth et al, *in prep.*). Contact NatureServe to learn more.
3. More information about the data and methods used to characterize resilience can be found at: http://bit.ly/TNC_resilience
4. Climate Science Special Report. The Fourth National Climate Assessment. U.S. Global Change Research Program. <https://science2017.globalchange.gov/>
5. U.S. Climate Adaptive capacity Toolkit, NOAA. <https://toolkit.climate.gov/>
6. National Park Service Climate Change Response Program: <https://www.nps.gov/orgs/ccrp/index.htm>
7. NatureServe, Climate Change Program <http://www.natureserve.org/biodiversity-science/conservation-topics/climate-change>
8. National Capital Region, Inventory & Monitoring Network: <https://science.nature.nps.gov/im/units/ncrn/>
9. National Park Service, National Capital Region, Natural Resources and Science, Urban Ecology Research Learning Alliance: <https://www.nps.gov/rlc/urbanecology/index.htm>

GLOSSARY OF TERMS

| | |
|------------------------------|--|
| Climate Change | Changes in weather patterns over relatively long time-scales. In this study, climate is defined based on averages and variability in temperature and precipitation for about 30-year periods. |
| Ecological System | A standardized ecological unit representing plant communities influenced by similar physical environments and dynamic ecological processes (like fire or flooding). Altered and disturbed vegetation are excluded. |
| Ecoregion | Provinces of unique physiography, soils, climate, and vegetation, containing geographically distinct assemblages of natural communities and species. |
| Enduring Features | The physical settings defined by landform, bedrock, soil, and topography are largely unchanged through time and provide the physical underpinnings for ecological diversity. |
| Landscape Diversity | Complex topography and elevation gradients creating a range of local temperature and moisture conditions, called micro-climates, within a given area. |
| Local Connectedness | The degree to which land cover patterns (e.g., agriculture, forest, wetlands) provide natural connections, supporting important ecological processes and the movement and dispersal of species. |
| Natural Habitats | An ecological area supporting native species including forests, wetlands, and native grasslands, but excluding areas with extensive human influence (e.g., regularly mowed meadows). |
| Novel | In terms of climate change exposure, "novel" refers to conditions that are higher or lower than the historical range of temperature or precipitation conditions observed in the past. |
| Non-Climate Stressors | External factors, unrelated to climate change, putting species and ecosystems at risk, such as invasive species, land use changes, predation, and disease. |
| Refugia | An area where environmental conditions allow a species or community to persist, even as unfavorable changes cause it to become extinct from surrounding areas. |
| Typical | In terms of climate change exposure, "typical" refers to conditions that are close to those experienced in the baseline past. |
| Species Turnover | Change in the types of plants and animals present at a site as new species move in and others are lost. |

REFERENCES

- Anderson, M.G., A. Barnett, M. Clark, C. Ferree, A. Olivero Sheldon, and J. Prince. 2016. Resilient Sites for Terrestrial Conservation in Eastern North America. The Nature Conservancy, Eastern Conservation Science.
- Daly, C., M. Halbleib, J.I. Smith, W.P. Gibson, M.K. Doggett, G.H. Taylor, J. Curtis, and P.P. Pasteris. 2008. Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *International Journal of Climatology*. 28(15), pp. 2031-2064.
- IPCC. 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- Oyler, J.W., A. Ballantyne, K. Jencso, M. Sweet, and S.W. Running. 2014. Creating a topoclimatic daily air temperature dataset for the conterminous United States using homogenized station data and remotely sensed land skin temperature. *International Journal of Climatology*, <http://dx.doi.org/10.1002/joc.4127>
- Stein, B.A., P. Glick, N. Edelson, and A. Staudt (eds.). 2014. Climate-Smart Conservation: Putting Adaptation Principles into Practice. National Wildlife Federation, Washington, D.C. www.nwf.org/ClimateSmartGuide
- Taylor, K. E., R.J. Stouffer, and G.A. Meehl. 2012. An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*. 93(4), 485-498.
- Full documentation of technical methods will be made available in a project report to be posted on the NPS IRMA portal pending project completion. For more information, contact Regan Smyth, NatureServe. Regan_Smyth@natureserve.org

Acknowledgments

Produced by NatureServe for the National Capital Region with the support of the National Park Service Climate Change Response Program. Special thanks for reviews by the National Park Service.

Suggested citation: Smyth, Regan, Judy Teague, Stephanie Auer, Allison Kenlan, and Healy Hamilton, 2017. Climate Change Vulnerability of Terrestrial Areas, Chesapeake & Ohio Canal National Historic Park. Unpublished Resource Brief, Department of the Interior, National Park Service. 10 p.